

**Angle-resolved photoemission study of heavy-fermion cerium
compounds CeRhIn₅ and CeIrIn₅**

Shin-ichi Fujimori¹, Akihiro Ino¹, Tetsuo Okane¹, Atushi Fujimori^{1,2}, Hisatomo Harima³,
Dai Aoki⁴, Yoshihumi Tokiwa⁵, Shuugo Ikeda⁵, Hiroaki Shishido⁵, Yoshinori Haga⁶,
Yoshichika Ōnuki^{5, 6}

¹ *Synchrotron Radiation Research Center, Japan Atomic Energy Research Institute,
SPring-8, Mikazuki, Hyogo 679-5148, Japan*

² *Department of Physics, Graduate School of Science, University of Tokyo, Bunkyo-ku,
Tokyo 113-0033, Japan*

³ *Institute of Scientific and Industrial Research, Osaka University, Osaka 567-0043,
Japan*

⁴ *DRFMC-SPSMS, CEA, 38054 Grenoble Cedex 9, France*

⁵ *Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan*

⁶ *Advanced Science Research Center, Japan Atomic Energy Research Institute, Tokai,
Ibaraki 319-1195, Japan*

Recently, two new ternary Ce-based heavy-fermion compounds CeRhIn₅ and CeIrIn₅ were synthesized. Both of them have tetragonal HoCoGa₅ structure, and their electronic structures are considered to be quasi-two-dimensional. Especially, it is reported that CeRhIn₅ is a pressure-induced superconductor*. In the present study, the electronic states of these two compounds are investigated by high-resolution angle-resolved photoelectron spectroscopy. The results are compared with the results of the FLAPW band structure calculations.

*H. Hagger *et al.*, Phys. Rev. Lett. **84**, 4986 (2000).